041

The A. & M. College of Texas.

DEPARTMENT OF OCEANOGRAPHY

OCEANOGRAPHIC SURVEY OF THE
GULF OF MEXICO

A CONTRIBUTION TO THE THEORY OF UPWELLING, PART II

Koji Hidaka

Research Conducted through the

Texas A. & M. Research Foundation

COLLEGE STATION, TEXAS

THE AGRICULTURAL AND MECHANICAL COLLEGE OF TEXAS
Department of Oceanography
College Station, Texas

Research conducted through the
Texas A. & M. Research Foundation

Project 24

A CONTRIBUTION TO THE THEORY OF UPWELLING, PART II
The Most Favorable Condition for Upwelling and
The Coastal Currents Induced by Winds

Dale F. Leipper
Project Supervisor

CONTENTS

TABLES

FIGURES

ii

A CONTRIBUTION TO THE THEORY OF UPWELLING. PART II
The Most Favorable Condition for Upwelling and
the Coastal Currents Induced by Winds

By

KOJI HIDAKA

ABSTRACT

In Technical Report No. 6 of this project the author showed the possibility of upwelling induced by a wind blowing parallel to the coast in such a manner that, in the northern (southern) hemisphere, the coast is on the left-hand (right-hand) side of an observer who looks in the direction of the wind. In this report he shows that an offshore wind blowing perpendicular to the coast can also give rise to a similar circulation in the same vertical plane, but in this case with much weaker intensity. From this result it can be concluded that the most intense upwelling will occur when the wind makes an angle of 21.5° with the coast line in an offshore direction. In addition to these circulations in the vertical plane there exist horizontal currents parallel to the coast. These will contribute to the formation of the coastal or longshore currents much debated recently. The horizontal components of the motion under consideration show a vertical variation very similar to the Ekman spiral at some distance from the coast.

1. **Upwelling Due to a Wind Not Parallel to the Coast.** In the first report on the subject of upwelling (Technical Report No. 6 of this project) the author has shown the possibility of upwelling induced by a wind blowing parallel to the coast. The result was that when a wind blows parallel to the coast in such a manner that, in the northern (southern) hemisphere an observer looking in the direction of the wind has the sea on his left-hand (right-hand) side, there exists an intense circulation in a vertical plane perpendicular to the coast ($x z$ -plane). This upwelling takes place close to the coast and has a velocity which is consistent with observations.

It may be anticipated, however, that a similar circulation can also be produced in the same vertical plane by an offshore wind, that is, a wind blowing offshore at right angles to the coast. To evaluate this circulation is, of course, possible by assuming a wind stress in the x- direction. Suppose we have a wind not perpendicular to, but making a certain angle with the coast, and divide the stress τ into its x- and y- components (Figure 1). Then the surface conditions (17) in Technical Report No. 6 should be replaced by

$$z = 0, \quad -A\frac{\partial u}{\partial z} = \tau, \quad -A\frac{\partial v}{\partial z} = \gamma,$$ (.)

where u and v are the x- and y- components of current velocity, τ_x and τ_y are those of wind stress while the dynamical and continuity equations and boundary conditions remain the same.

In exactly the same manner as in the first report, we can integrate the equations of motion and obtain the stream function in the vertical plane perpendicular to the coast and the components u and v of the horizontal current. They are

$$\psi(x,z) = \text{real part of}$$

$$\frac{2\pi(\tau_x + i\tau_y)}{\rho\,w\sin\varphi}\int_0^\infty \frac{\sin\lambda\frac{x}{D_r}\left(1-\sin\lambda\frac{L}{D_h}\right)}{\lambda}\cdot\frac{e^{-\sqrt{\lambda^2+2\pi^2 i}\,\frac{z}{D_r}}-1}{\lambda^2+2\pi^2 i}\,d\lambda \qquad (2)$$

and

$$u+iv = \frac{2\pi(\tau_x+i\tau_y)}{\rho\,w\sin\varphi\,D_v}\int_0^\infty \frac{\sin\frac{x}{D_h}\left(1-\sin\lambda\frac{L}{D_h}\right)}{\lambda}\cdot\frac{e^{-\sqrt{\lambda^2+2\pi^2 i}\,\frac{z}{D_v}}}{\sqrt{\lambda^2+2\pi^2 i}}\,d\lambda \qquad (3)$$

respectively. These expressions can be put in real forms,

$$\psi(x,z) = \frac{2\pi\,\tau_x}{\rho\,w\sin\varphi}\,\Phi_x(x,z) + \frac{2\pi\,\tau_y}{\rho\,w\sin\varphi}\,\Phi_y(x,z)\;, \qquad (4)$$

$$u = \frac{2\pi\,\tau_x}{\rho\,w\sin\varphi\,D_v}\,N(x,z) - \frac{2\pi\,\tau_y}{\rho\,w\sin\varphi\,D_v}\,M(x,z)\;, \qquad (5)$$

$$v = \frac{2\pi\,\tau_x}{\rho\,w\sin\varphi\,D_v}\,M(x,z) + \frac{2\pi\,\tau_y}{\rho\,w\sin\varphi\,D_v}\,N(x,z)\;, \qquad (6)$$

where

$$\Phi_x(x,z) = \int_0^\infty \frac{(P^2-Q^2)(e^{-P\frac{z}{D_r}}\cos Q\frac{z}{D_r}-1)-2PQ\,e^{-P\frac{z}{D_r}}\sin Q\frac{z}{D_r}}{(P^2+Q^2)^2}\;\frac{\sin\lambda\frac{x}{D_h}\left(1-\cos\lambda\frac{L}{D_h}\right)}{\lambda}\,d\lambda\;, \qquad (7)$$

$$\Phi_y(x,z) = \int_0^\infty \frac{(P^2-Q^2)\,e^{-P\frac{z}{D_r}}\sin Q\frac{z}{D_r}+2PQ(e^{-P\frac{z}{D_v}}\cos Q\frac{z}{D_v}-1)}{(P^2+Q^2)^2}\;\frac{\sin\lambda\frac{x}{D_h}\left(1-\cos\lambda\frac{L}{D_h}\right)}{\lambda}\,d\lambda \qquad (8)$$

$$M(x,z) = \int_0^\infty \frac{Q\cos Q\frac{z}{D_v}+P\sin Q\frac{z}{D_v}}{P^2+Q^2}\,e^{-P\frac{z}{D_v}}\;\frac{\sin\lambda\frac{x}{D_h}\left(1-\cos\lambda\frac{L}{D_h}\right)}{\lambda}\,d\lambda \qquad (9)$$

$$N(x,z) = \int_0^\infty \frac{P_{\min} Q \frac{z}{D_r} - Q_{\min} Q \frac{z}{D_r}}{P^2 + Q^2} e^{-P\frac{x}{A}} \; \operatorname{cod} \frac{z}{D_h}\left(1 - \operatorname{cod}\frac{L}{D_h}\right) d\lambda , \quad (10)$$

and we have

$$D_h = \pi \sqrt{A_h / \rho \omega \sin \phi} , \qquad (11)$$

$$D_v = \pi \sqrt{A_v / \rho \omega \sin \phi} . \qquad (12)$$

Here A_h and A_v are the coefficients of horizontal and vertical mixing, ρ the density of sea water, ω the angular velocity of the Earth's rotation and ϕ the geographic latitude.

If we put

$$\tau_x = 0 , \quad \tau_y = \tau ,$$

we have exactly the problem we discussed in the first report, in which the wind blows parallel to the coast.

The stream function $\phi_x(x,z)$ and $\phi_y(x,z)$, and the function $M(x,z)$ and $N(x,z)$ which give the horizontal components of velocity were evaluated by numerical computation against several values of x/D_h and z/D_v, and are compiled in Tables I to IV.

Since a wind of any direction can be divided into its two horizontal components perpendicular and parallel to the coast, it will be sufficient for us to discuss only the vertical circulations induced by the offshore and longshore winds and to combine the result.

2. Pattern of the Vertical Circulations. Figures 2 and 3 show the stream lines in the vertical circulations induced in the plane perpendicular to the coast (xz — plane) by longshore and offshore winds respectively. There seems to be an essential difference between them. Both are clockwise circulations and the stream functions have negative values. In each of these two cases, the water is upwelled from levels deeper than $z = D_v$, but, as a whole, it appears that a longshore wind originates upwelled water from deeper layers than does an offshore wind. The circulation induced by a longshore wind has already been described in detail in the first report. The circulation due to an offshore wind of the same stress carries a much smaller amount of water than does that due to a longshore wind. Moreover, the former has a rather complicated structure, having two eddies (Figure 3) in upper layers, one situated close to the coast

and the other near the outer boundary of the wind belt. This means that upwelling due to a longshore wind is far more effective in lowering the surface temperature of the coastal regions than that induced by an offshore one, since the former carries much larger amounts of colder water to the surface from deeper layers than the latter.

3. __Most Favorable Condition for Upwelling.__ The amount T of water upwelled across a long horizontal strip 1 cm wide extending from the coast to a distance x can be evaluated as

$$T = \int_0^x \rho \omega \, dx = \rho \int_0^x \frac{\partial \psi}{\partial x} \, dx$$

$$= \rho \, \psi(x, z) - \rho \, \psi(0, z)$$

$$= \rho \, \psi(x, z) , \qquad\qquad (13)$$

since the stream function vanishes along the coastal wall ($x = 0$). Thus the vertical mass transport T will be largest when we take x at a distance at which the functions ϕ_x and ϕ_y have the largest negative value. By inspection of Tables I and II and Figures 2 and 3, we can estimate

$$T_y = -0.090 \times \frac{2\pi \, \bar{\tau}_y}{\omega \sin \varphi} \qquad \text{by longshore wind} \qquad (14)$$

and

$$T_x = -0.0355 \times \frac{2\pi \, \bar{\tau}_x}{\omega \sin \varphi} \qquad \text{by offshore wind} \qquad (15)$$

If the wind blows in a direction making an angle θ with the coast in an offshore direction (Figure 1), we have

$$\bar{\tau}_x = \bar{\tau} \sin \theta , \quad \bar{\tau}_y = \bar{\tau} \cos \theta \qquad\qquad (16)$$

where $\bar{\tau}$ is the absolute magnitude of the wind stress. Then we have from (14) and (15)

$$T = T_x + T_y$$

$$= (-0.0355 \sin \theta - 0.090 \cos \theta) \frac{2\pi \, \bar{\tau}}{\omega \sin \varphi} . \qquad (17)$$

This gives an approximate amount of water upwelled to the surface layers per unit length of the coast. The most intense upwelling will therefore take place when $d\overline{I}/d\theta = 0$, or

$$ \tan \theta = \frac{0.0355}{0.090} \quad \text{or} \quad \theta = 21.5° \tag{18} $$

providing the magnitude of wind stress $\overline{\zeta}$ remains the same. This means that the upwelling will be most intense when the wind makes an angle of 21.5° with the coast line in an offshore direction, that is, when it is slightly deviated offshore from the coast line. Of course, we have more or less upwelling in a sector within ± 90° of this direction. According to Dr. Arnold Glaser, upwelling off the coast of Lima, Peru, appears to be more intense when the southerlies blow slightly deviated in an offshore direction rather than when they blow parallel to the coast.

In the Gulf of Mexico, we have easterlies almost all the year round. This means that we can always expect upwelling along the northern coast of Cuba and Yucatan coast of Mexico. We do not have any distinct evidence of upwelling and the existence of colder water off these coasts. But we know that many American anglers go to fish in the waters off the Yucatan coast and special boats are sent from the United States coasts for these people. This fact may suggest the existence of upwelling in this region. According to the surface observations of the Department of Oceanography, Agricultural and Mechanical College of Texas made recently, there are also some indications of colder surface temperatures in the western part of the Gulf of Mexico very close to the coast. If they really exist, this temperature distribution may be ascribed to upwelling due to the easterly winds prevailing in these latitudes.

4. Coastal Currents. Recently several observations (Putnam, Munk and Traylor, 1949; Shepard and Inman, 1950, 1951; Inman and Quinn, 1953) on the flow of water close or adjacent to coasts have been reported. Many authors have discussed this subject, but they have not yet been able to arrive at a satisfactory explanation of possible causes of coastal or longshore currents. However, most of these currents are ascribed to the action of surf and waves. The longshore component which we have obtained may explain some part of these flows. The expression for this component is given by

$$ v = \frac{2\pi \overline{\zeta}_x}{\rho w \sin \phi D_r} N(\chi, z) - \frac{2\pi \overline{\zeta}_y}{\rho w \sin \phi D_r} M(\chi, z) \tag{19} $$

where M and N are the functions compiled in Tables III and IV and illustrated in Figures 4 and 5. The two terms in the right-hand member of (19) show the longshore currents induced by offshore and longshore winds respectively. If the wind is parallel to the coast we have $\overline{\zeta}_x = 0$ and

TABLE I.   VALUES OF THE STREAM FUNCTION

$$\Phi_y(x, z) \times 10^4$$

z/D_c \ x/D_h	0	.1396	.2793	.5585	.8378	1.1170	1.3963	1.6755	1.9548
0	0	0	0	0	0	0	0	0	0
.2	0	-175	-313	-449	-461	-465	-455	-425	-310
.4	0	-277	-499	-725	-748	-750	-734	-682	-496
.6	0	-321	-578	-843	-867	-866	-848	-788	-576
.8	0	-334	-600	-871	-891	-887	-869	-808	-589
1.0	0	-332	-598	-865	-890	-873	-856	-797	-579
1.2	0	-329	-592	-852	-863	-853	-837	-782	-568
1.4	0	-327	-587	-844	-853	-843	-827	-773	-561
1.6	0	-326	-585	-840	-848	-837	-823	-769	-558

TABLE II.   VALUES OF STREAM FUNCTION

$$\Phi_y(x, z) \times 10^4$$

z/D_c \ x/D_h	0	.1396	.2793	.5585	.8378	1.1170	1.3963	1.6755	1.9548
0	0	0	0	0	0	0	0	0	0
.2	0	-168	-303	-301	-254	-270	-280	-305	-202
.4	0	-194	-342	-310	-230	-238	-257	-303	-199
.6	0	-179	-309	-234	-135	-135	-161	-226	-133
.8	0	-162	-275	-172	- 68	= 58	- 83	-162	- 85
1.0	0	-152	-256	-138	- 19	- 17	- 43	-129	- 61
1.2	0	-148	-248	-125	- 5	- 3	- 30	-107	- 54
1.4	0	-148	-274	-124	- 4	- 3	- 30	-107	- 54
1.6	0	-147	-248	-126	- 6	- 6	- 33	-110	- 57

TABLE III. VALUES OF THE FUNCTION

$$N(x,z) \times 10^3$$

z/Dv \ x/Dh	0	.1396	.2793	.5585	.8378	1.1170	1.3963	1.6755	1.9548
0	0	+240	+277	+284	+253	+252	+252	+266	+243
.2	0	+ 41	+ 64	+ 55	+ 35	+ 31	+ 38	+ 47	+ 22
.4	0	- 3	- 8	- 22	- 42	- 45	- 42	- 20	- 17
.6	0	- 10	- 20	- 38	- 47	- 50	- 48	- 39	- 29
.8	0	- 7	- 14	- 24	- 29	- 30	- 29	- 24	- 20
1.0	0	- 3	- 6	- 11	- 12	- 12	- 12	- 10	- 8
1.2	0	- 1	- 2	- 2	- 3	- 3	- 2	- 2	- 1
1.4	0	0	0	- 1	+ 1	+ 1	+ 1	+ 1	+ 1
1.6	0	0	+ 1	+ 1	+ 2	+ 2	+ 2	+ 1	+ 1

TABLE IV. VALUES OF THE FUNCTION

$$M(x,z) \times 10^3$$

z/Dv \ x/Dh	0	.1396	.2793	.5585	.8378	1.1170	1.3963	1.6755	1.9548
0	0	- 98	-174	-248	-255	-243	-263	-235	-174
.2	0	- 70	-127	-186	-191	-182	-197	-175	-129
.4	0	- 34	- 62	- 93	- 96	- 90	- 97	- 85	- 62
.6	0	- 12	- 21	- 31	- 30	- 26	- 30	- 26	- 18
.8	0	- 2	- 3	- 2	0	+ 3	+ 1	+ 1	+ 1
1.0	0	+ 1	+ 3	+ 6	+ 10	+ 10	+ 9	+ 8	+ 6
1.2	0	+ 2	+ 3	+ 6	+ 8	+ 8	+ 8	+ 6	+ 5
1.4	0	+ 1	+ 2	+ 3	+ 4	+ 4	+ 4	+ 3	+ 2
1.6	0	0	+ 1	+ 1	+ 1	+ 1	+ 1	+ 1	+ 1

$$v = -\frac{2\pi\,\tau_y}{\rho\omega\sin\phi\,D_v}\,M(\chi,z),\tag{20}$$

and if the wind blows in an offshore direction perpendicular to the coast we have

$$v = +\frac{2\pi\,\tau_x}{\rho\omega\sin\phi\,D_v}\,N(\chi,z).\tag{21}$$

From the Tables III and IV and Figures 4 and 5 it is possible for us to tell the difference between the two cases. The coastal current induced by a longshore wind is directed leeward as far down as the level $z = 0.9\,D_v$, but the direction of flow is reversed at deeper levels. On the other hand, the longshore current produced by an offshore wind flows to the right of the wind direction only in layers shallower than $z = 0.3\,D_v$. There is a secondary maximum of velocity in the opposite direction at the approximate level $z = 0.5\,D_v$.

5. **Vertical Variation of Horizontal Currents: Ekman Spiral.** The expressions

$$
\begin{aligned}
u &= \frac{2\pi\,\tau_x}{\rho\omega\sin\phi\,D_v}\,N(\chi,z) - \frac{2\pi\,\tau_y}{\rho\omega\sin\phi\,D_v}\,M(\chi,z), \\[2mm]
v &= \frac{2\pi\,\tau_x}{\rho\omega\sin\phi\,D_v}\,M(\chi,z) - \frac{2\pi\,\tau_y}{\rho\omega\sin\phi\,D_v}\,N(\chi,z),
\end{aligned}
\tag{22}
$$

for the horizontal currents show that the angle between the wind and current induced at a given depth depends only upon the distance from the coast and not on the direction of wind. The angle between the wind and the surface current induced by it is computed for several points at different distances from the coast and compiled in Table V.

Table V

The Angle Between the Wind and Surface Current

x/D_v	0.0000	.1396	.2793	.5585	.8378	1.1170	1.3963	1.6755	1.9548
Angle	0°	22.2°	32.2°	41.2°	45.2°	46.3°	46.3°	41.5°	35.7°

Thus the angle between the wind and surface current is nearly 45° at the middle part of the wind belt, but decreases both toward the coast and outer margin of the wind belt. Along the coast there is a very weak current in approximately the same direction as the wind. The vertical variation of currents is practically equal to the Ekman spiral along the

median line of the wind belt, but is more or less flattened close to the
coast and near the outer margin of the wind belt. Some of these spirals
are illustrated in Figure 6.

References

Inman, D. L. and W. R. Quinn, 1953. Currents in the surf zone. Coastal
    Engineering. Chapter 3, p. 24 - 36. (Contributions from the Scripps
    Institution of Oceanography No. 560)

Putnam, J. A., W. H. Munk and M. A. Traylor, 1949. The prediction of
    longshore currents. Ameri. Geophys. Union Trans. Vol. 30, p. 337 - 345.

Shepard, F. P. and D. L. Inman, 1950. Nearshore water circulation related
    to bottom topography and wave refraction. Ameri. Geophys. Union Trans.
    Vol. 31, p. 196 - 212.

_____, 1950. Nearshore circulation, Proceedings of the 1st Conference
    on Coastal Engineering, Council on Wave Research, Engineering Founda-
   tion. p. 50 - 59.

FIGURE 1

WIND MAKING AN ANGLE
θ WITH THE COAST

UPWELLING AS INDUCED BY A WIND PARALLEL TO THE COAST
(ILLUSTRATED BY THE STREAMLINES IN THE VERTICAL PLANE PERPENDICULAR TO THE COAST)

FIGURE 2

FIGURE 3

UPWELLING AS INDUCED BY AN OFFSHORE WIND

(illustrated by the stream-lines in the vertical plane perpendicular to the coast)

FIGURE 4

DISTRIBUTION OF LONGSHORE CURRENTS M(x,z)
INDUCED BY A WIND τ_x PARALLEL TO THE COAST

FIGURE 5

DISTRIBUTION OF LONGSHORE CURRENT N(x,z)
INDUCED BY AN OFFSHORE WIND τ_y

FIGURE 9
VERTICAL VARIATION OF DRIFT CURRENTS

No. Copies	Address
8	Chief of Naval Research Navy Department Washington 25, D. C. Attn: Code 416 (2) " Code 446 (3) " Code 466 (3)
1	ONR Resident Representative University of Texas Main Building, Room 2506 Austin 12, Texas
1	Director Office of Naval Research 150 Causeway Street Boston 14, Massachusetts
1	Director Office of Naval Research 1000 Geary Street San Francisco 9, California
1	Director Office of Naval Research Tenth Floor, 86 E. Randolph St. Chicago 1, Illinois
1	Director Office of Naval Research 346 Broadway New York 13, New York
1	Director Office of Naval Research 1030 E. Green Street Pasadena 1, California
5	U. S. Navy Hydrographic Office Washington 25, D. C. Attn: Division of Oceanography
5	Chief, Bureau of Ships Navy Department Washington 25, D. C. Attn: Code 847 (2) " Code 516 (2) " Code 845 (1)
4	Director U. S. Naval Electronics Lab. San Diego 52, California Attn: Code 550, 551, 552, 610
1	Commanding Officer U. S. Navy Underwater Sound Lab. New London, Connecticut
9	Naval Research Laboratory Technical Services Washington 25, D. C.
1	Chief, Bureau of Yards & Docks Navy Department Washington 25, D. C.
1	Commanding Officer Naval Ordnance Laboratory White Oak Silver Spring 19, Maryland
2	Asst. Naval Attache for Research American Embassy Navy 100 Fleet Post Office, New York
3	British Joint Services Mission Main Navy Building Washington 25, D. C.
1	Commanding General Research and Development Div. Department of the Army Washington 25, D. C.
1	U. S. Army Beach Erosion Board 5201 Little Falls Road, NW Washington 16, D. C.
1	The Chief, Armed Forces Special Weapons Project P. O. Box 2610 Washington, D. C.
1	Office of the Quartermaster General Military Planning Division Research and Development Branch Washington 25, D. C.
1	Research & Development Board National Military Establishment Washington 25, D. C. Attn: Committee on Geophysics and Geography

No. Copies	Address
1	U. S. Weather Bureau U. S. Department of Commerce Washington 25, D. C. Attn: Scientific Services
1	Commanding Officer Cambridge Field Station 230 Albany Street Cambridge 39, Massachusetts Attn: CRHSL
1	Commandant (OAO) U. S. Coast Guard Washington 25, D. C.
1	Director U. S. Coast and Geodetic Survey Department of Commerce Washington 25, D. C.
1	U. S. Fish & Wildlife Service P. O. Box 3830 Honolulu, T. H.
1	U. S. Fish & Wildlife Service Woods Hole, Massachusetts
1	U. S. Fish & Wildlife Service South Atlantic Offshore Fishery Investigations c/o Georgia Game & Fish Commission P. O. Box 312 Brunswick, Georgia
2	U. S. Fish & Wildlife Service Fort Crockett Galveston, Texas
3	Director U. S. Fish & Wildlife Service Department of the Interior Washington 25, D. C. Attn: Dr. L. A. Walford
1	U. S. Fish & Wildlife Service 450 B. Jordan Hall Stanford University, California
1	Secretary Gulf States Marine Fisheries Commission Audubon Building 931 Canal Street New Orleans 16, Louisiana
1	Secretary Atlantic States Marine Fisheries Commission 22 West 1st Street Mount Vernon, New York Attn: Wayne D. Heydecker
1	Texas Game and Fish Commission Rockport, Texas
2	California Academy of Sciences Golden Gate Park San Francisco, California Attn: Dr. R. C. Miller
1	Director Institute of Marine Science Port Aransas, Texas
1	Director Gulf Coast Research Laboratory Ocean Springs, Mississippi
1	Director Narragansett Marine Laboratory Kingston, Rhode Island
1	Director, Hawaii Marine Laboratory University of Hawaii Honolulu, T. H.
1	Director, Marine Laboratory University of Miami Coral Gables, Florida
1	American Museum of Natural History Central Park West at 79th Street New York 24, New York
1	Director Lamont Geological Observatory Torrey Cliff Palisades, New York
1	Department of Conservation Cornell University Ithaca, New York Attn: Dr. J. Ayers

No. Copies	Address
1	Department of Zoology Rutgers University New Brunswick, New Jersey Attn: Dr. H. K. Haskins
2	Scripps Institution of Oceanography La Jolla, California Attn: Director (1) " Library (1)
1	The Oceanographic Institute Florida State University Tallahassee, Florida
1	Department of Engineering, Dean University of California Berkeley, California
2	Director Woods Hole Oceanographic Institute Woods Hole, Massachusetts
1	Director Chesapeake Bay Institute Box 426A RFD 2 Annapolis, Maryland
2	Head, Department of Oceanography University of Washington Seattle, Washington
1	Bingham Oceanographic Foundation Yale University New Haven, Connecticut
1	Allan Hancock Foundation University of Southern California Los Angeles 7, California
1	Head, Department of Biology Texas Christian University Ft. Worth 9, Texas
1	Head, Dept. Meteorology & Ocean. New York University New York, New York
1	Bear's Bluff Laboratory Wadmalaw Island, South Carolina Attn: Dr. Robert Lunz
1	Department of Biology University of Florida Gainesville, Florida Attn: Mr. H. Love Pierce
1	Director, Virginia Fisheries Labor. College of William and Mary Gloucester Point, Virginia
1	Director, University of Florida Marine Biological Station Gainesville, Florida
1	Director, Alabama Marine Laboratory Bayou La Batre, Alabama
1	Southern Regional Education Board Marine Sciences 830 West Peachtree Street, N. W. Atlanta, Georgia
1	Director, Louisiana State University Marine Laboratory Baton Rouge, Louisiana
1	Director, Institute of Fisheries Research University of North Carolina Morehead City, North Carolina
1	Director Duke University Marine Laboratory Beaufort, North Carolina
1	Institute of Engineering Research 214 Naces Hall Berkeley 4, California Attn: Prof. J. W. Johnson
1	Chief, Air Weather Service Department of the Air Force Washington, D. C.
1	Head, Department of Oceanography Brown University Providence, Rhode Island